arXiv:1603.08945v2 [cond-mat.stat-mech] 30 May 2016

On two-time distribution functions in (1+1) random directed polymers

Victor Dotsenko
LPTMC, Université Paris VI, Paris, France and
L.D. Landau Institute for Theoretical Physics, Moscow, Russia
(Dated: May 31, 2016)

An asymptotic analytic expression for the two-time free energy distribution function in (1+1) random directed polymers is derived in the limit when the two times are close to each other.

PACS numbers: 05.20.-y 75.10.Nr 74.25.Qt 61.41.+e

In this paper we consider $(1+1)$ directed polymers model in terms of a scalar field $\phi(\tau)$ within an interval $0 \le \tau \le t$ defined by the Hamiltonian

$$H[\phi(\tau), V] = \int_0^t d\tau \Big\{ \frac{1}{2} \big[\partial_\tau \phi(\tau)\big]^2 + V[\phi(\tau), \tau] \Big\}; \tag{1}$$

where $V(\phi, \tau)$ is the Gaussian distributed random potential with a zero mean, $\overline{V(\phi,\tau)} = 0$, and δ-correlations, $\overline{V(\phi,\tau)V(\phi',\tau')} = u\delta(\tau-\tau')\delta(\phi-\phi')$. The parameter u is the strength of the disorder. This problem, which is equivalent to the one of the KPZ-equation [1] describing the time evolution of an interface in the presence of noise, has been the focus of intense studies during past three decades [2–26]. At present it is well established that the fluctuations of the free energy of the model defined by the Hamiltonian (1) are described by the Tracy-Widom distribution [27]. The two-point free energy distribution function which describes joint statistics of the free energies of the directed polymers coming to two different endpoints has been derived in [28–30]. The generalization for the corresponding N-point distribution has been obtained in [31, 32].

All the above studies were devoted to the so called "one-time" situation. The joint probability distribution function at two different times has been studied in the paper [33, 34] (in terms of non-rigorous replica Bethe ansatz approach) and in the paper [35] (mathematically rigorous derivation). Unfortunately, the results obtained in these papers are expressed in terms rather complicated mathematical objects whose analytic properties are not clear. Quite recently the time correlations of the KPZ interface growth has been studied both theoretically, in terms of the variational problem involving two independent Airy processes [36], and experimentally [37].

In this paper I am going to consider the two-time free energy distribution functions of the directed polymers defined by the Hamiltonian (1) in the limit when the two times are getting close to each other. For the fixed boundary conditions the partition function of this model is

$$Z(x;\, t) = \int_{\phi(0)=0}^{\phi(t)=x} \mathcal{D}\phi(\tau)\; \mathrm{e}^{-\beta H[\phi]} \;=\; \exp\big\{-\beta F(x;t)\big\} \tag{2}$$

where β is the inverse temperature and $F(x,t)$ is the free energy. In the limit $t \to \infty$ the free energy scales as

$$\beta F(x,t) = \beta f_0 t + \beta x^2/2t + \lambda(t) f(x)\ , \tag{3}$$

where f_0 is a non-random free energy density,

$$\lambda(t) = \frac{1}{2}(\beta^5 u^2 t)^{1/3} \propto t^{1/3} \tag{4}$$

and $f(x)$ is a random quantity described by the Tracy-Widom distribution. As the first two trivial terms of the free energy, eq.(3), can be easily eliminated by simple redefinition of the partition function, they will be omitted in the further calculations, in other words,

$$Z(x;\, t) = \exp\big\{-\lambda(t) f(x)\big\}\ . \tag{5}$$

Let us consider two directed polymer trajectories with the fixed boundary conditions at two different times: $\phi_1(0) = \phi_1(t) = 0$ and $\phi_2(0) = \phi_2\big[(1-\Delta)t\big] = 0$ having the free energies f_1 and f_2 correspondingly. By definition

$$\exp\big\{-\lambda(t) f_1\big\} \;=\; Z(0;\, t) \tag{6}$$

and

$$\exp\big\{-\lambda\big[(1-\Delta)t\big] f_2\big\} \;=\; Z(0;\, t-\Delta t) \tag{7}$$

where the partition function $Z(x;t)$ is defined in eq.(2). We are going to study the properties of the joint probability distribution function, $W(f_1, f_2, \Delta)$, the limit $t \to \infty$ when the parameter $0 < \Delta < 1$ remains finite.

By definition, the probability distribution function $W(f_1, f_2, \Delta)$ is related with the corresponding probability density $P_{t,\Delta}(f_1, f_2)$ as

$$W(f_1, f_2, \Delta) \;=\; \lim_{t\to\infty} \int_{f_1}^{+\infty} df_1' \int_{f_2}^{+\infty} df_2' \; P_{t,\Delta}(f_1', f_2') \;=\; \int_{f_1}^{+\infty} df_1' \int_{f_2}^{+\infty} df_2' \; \mathcal{P}_{\Delta}(f_1', f_2') \tag{8}$$

where

$$\mathcal{P}_{\Delta}(f_1, f_2) \;=\; \lim_{t\to\infty} P_{t,\Delta}(f_1, f_2)\,. \tag{9}$$

In terms of the above partition functions (6)-(7) the probability distribution function $W(f_1, f_2, \Delta)$ can be defined as follows:

$$W(f_1, f_2, \Delta) = \lim_{t\to\infty} \sum_{N,K=0}^{\infty} \frac{(-1)^{N+K}}{N!\,K!} \exp\big(\lambda_t N f_1 + \lambda_t K f_2\big)\; \overline{Z^N(0;t)\, Z^K(0; t-\Delta t)} \tag{10}$$

where $\overline{(...)}$ denotes the disorder averaging and the parameter $\lambda(t)$ is defined in eq.(4). Indeed, substituting here eqs.(6) and (7) we get

$$\begin{aligned} W(f_1, f_2, \Delta) \;&=\; \lim_{t\to\infty} \sum_{N,K=0}^{\infty} \frac{(-1)^{N+K}}{N!\,K!} \int_{-\infty}^{+\infty} df_1' \int_{-\infty}^{+\infty} df_2' \; P_{t,\Delta}(f_1', f_2') \exp\Big\{\lambda(t) N (f_1 - f_1') + \lambda[(1-\Delta)t] K (f_2 - f_2')\Big\} \\ &=\; \lim_{t\to\infty} \int_{-\infty}^{+\infty} df_1' \int_{-\infty}^{+\infty} df_2' \; P_{t,\Delta}(f_1', f_2') \exp\Big\{ - \exp\big\{\lambda(t)(f_1 - f_1')\big\} - \exp\big\{\lambda[(1-\Delta)t](f_2 - f_2')\big\}\Big\} \\ &=\; \int_{-\infty}^{+\infty} df_1' \int_{-\infty}^{+\infty} df_2' \; \mathcal{P}_{\Delta}(f_1', f_2')\; \theta\big(f_1' - f_1\big)\theta\big(f_2' - f_2\big) \end{aligned} \tag{11}$$

which coincides with the definition, eq.(8).

According to the definition of the partition function, eq.(2),

$$Z(0;\, t) \;=\; \int_{-\infty}^{+\infty} dx \; Z(x;\, t - \Delta t)\, Z^*(x;\, \Delta t) \tag{12}$$

where $Z^*(x;\, \Delta t)$ is the partition function of the directed polymer system in which time goes backwards, from t to $(t - \Delta t)$. Substituting eq.(12) into eq.(10) we get

$$\begin{aligned} W(f_1, f_2, \Delta) \;&=\; \lim_{t\to\infty} \sum_{N,K=0}^{\infty} \frac{(-1)^{N+K}}{N!\,K!} \exp\Big[\lambda(t) N f_1 + \lambda(t - \Delta t) K f_2\Big] \times \\ &\times \int_{-\infty}^{+\infty} dx_1 \,...\, dx_N \overline{\Big[\prod_{a=1}^{N} Z(x_a;\, t - \Delta t)\Big] Z^K(0;\, t - \Delta t) \Big[\prod_{a=1}^{N} Z^*(x_a;\, \Delta t)\Big]} \end{aligned} \tag{13}$$

or

$$\begin{aligned} W(f_1, f_2, \Delta) \;&=\; \lim_{t\to\infty} \sum_{N,K=0}^{\infty} \frac{(-1)^{N+K}}{N!\,K!} \exp\Big[\lambda(t) N f_1 + \lambda(t - \Delta t) K f_2\Big] \times \\ &\times \int_{-\infty}^{+\infty} dx_1 ... dx_N \Psi\big(x_1, ..., x_N, \underbrace{0, ..., 0}_{K};\; (t - \Delta t)\big)\; \Psi^*\big(x_1, ..., x_N;\; \Delta t\big) \end{aligned} \tag{14}$$

where

$$\Psi(x_1, ..., x_N;\; t) \;\equiv\; \overline{Z(x_1;\, t)\, Z(x_2\, t) \,...\, Z(x_N\, t)} \tag{15}$$

One can easily show that $\Psi(\mathbf{x};\,t)$ is the wave function of N-particle quantum boson system with attractive δ-interaction:

$$\beta\,\partial_t\Psi(\mathbf{x};t) = \frac{1}{2}\sum_{a=1}^{N}\partial_{x_a}^2\Psi(\mathbf{x};t) + \frac{1}{2}\,\beta^3 u\sum_{a\neq b}^{N}\delta(x_a-x_b)\;\Psi(\mathbf{x};t) \tag{16}$$

with the initial condition $\Psi(\mathbf{x};\,0)=\Pi_{a=1}^{N}\delta(x_a)$. The time dependent wave function $\Psi(\mathbf{x};\,t)$ of this quantum problem can be represented in terms of the linear combination of the Bethe ansatz eigenfunctions of eq.(16) which are well known since long time ago (for details see e.g.[38–42])

Using this representation the general (rather cumbersome) result for $W(f_1,f_2,\Delta)$, eq.(14) (for an arbitrary Δ) has been derived in [33] (see eqs(73), (76) and (79)-(82)). In this paper we would like to study the limit $\Delta\to 0$ which is somewhat tricky as it is considered *after* the limit $t\to\infty$ is already taken.

It is clear that at $\Delta=0$ the two free energies f_1 and f_2 must coincide, so that $\mathcal{P}_{\Delta=0}(f_1;f_2)\;=\;\delta(f_1-f_2)\,P_{GUE}(f_1)$, and $W(f_1,f_2,\Delta=0)\;=\;F_2(-\max\{f_1,f_2\})$ where $F_2(-f)$ is the GUE Tracy-Widom distribution. However, when $\Delta\ll 1$ but not zero, the situation becomes much less trivial. In this case the distribution density $P_\Delta(f_1;f_2)$ have a shape of a narrow "δ-like" peak with a width $\sim\Delta^{1/3}$ around f_1-f_2. Then, if $|f_1-f_2|\gg\Delta^{1/3}$, the "δ-like" distribution $P_\Delta(f_1;f_2)$ can be treated as the δ-function, so that one would get $W(f_1,f_2,\Delta)\simeq F_2(-\max\{f_1,f_2\})$ again. However, if $|f_1-f_2|\sim\Delta^{1/3}$ the situation becomes much more complicated.

Let us introduce a new scaling variable

$$\xi\;=\;\frac{f_1-f_2}{\Delta^{1/3}\,2^{2/3}} \tag{17}$$

where the factor $2^{2/3}$ is introduced just to simplify the final formula. Substituting $f_2=2^{2/3}f$ and $f_1=2^{2/3}f\;+\;\Delta^{1/3}\,2^{2/3}\,\xi$ into $W(f_1,f_2,\Delta)$ and taking the limit $\Delta\to 0$ (so that ξ remains finite) we define a new probability distribution function

$$W(f,\xi)\;\equiv\;\lim_{\Delta\to 0}W\Big(2^{2/3}f\;+\;2^{2/3}\Delta^{1/3}\,\xi,\;\;2^{2/3}f;\;\Delta\Big) \tag{18}$$

which describe the "detailed shape" of the distribution $W(f_1,f_2,\Delta)$ in the limit $\Delta\to 0$ when the two free energies f_1 and f_2 are close to each other $(f_1-f_2)\;\sim\;\Delta^{1/3}$.

It can be shown (see Appendix) that the result for $W(f,\xi)$ can be represented in terms of two Fredholm determinants in the form of the following rather compact analytic formula:

$$W(f,\xi)\;=\;\frac{1}{2\pi i}\int_{|z|=1}\frac{dz}{z}\det\Big[\hat{1}\;-\;(1-z)\hat{K}_{-f}\Big]\;\det\Big[\hat{1}\;+\;z^{-1}\hat{K}_{-\xi}\Big] \tag{19}$$

where the integration over z goes in the complex plane around zero and $\hat{K}_{-f}$ is the integral operator with the usual Airy kernel $K(u-f;u'-f)$ (with $(u,\,u')\;\geq 0$):

$$K(u-f;u'-f)\;=\;\int_0^{+\infty}dy\;\;\mathrm{Ai}(y+u-f)\;\;\mathrm{Ai}(y+u'-f) \tag{20}$$

Following the standard transformations of the Fredholm determinant with the Airy kernel [27] the distribution function, eq.(19), can equivalently be represented as

$$W(f,\xi)\;=\;\frac{1}{2\pi i}\int_{|z|=1}\frac{dz}{z}\exp\left\{-\int_{-f}^{\infty}dt\,(t+f)\;q_1^2(t)\;-\;\int_{-\xi}^{\infty}dt\,(t+\xi)\;q_2^2(t)\right\} \tag{21}$$

where $q_1(t)$ and $q_2(t)$ are the solutions of the Panlevé II equation

$$q''(t)=tq(t)+2q^3(t) \tag{22}$$

with the boundary conditions, $q_1(t\to+\infty)\sim(1-z)^{1/2}\;\mathrm{Ai}(t)$ and $q_2(t\to+\infty)\sim(-z)^{-1/2}\;\mathrm{Ai}(t)$ correspondingly. Eqs.(19)-(22) constitute the main result of the present paper. The analytic properties of the function $W(f,\xi)$ given by the above formulas are still to be investigated.

Acknowledgments

I am grateful to Alexei Borodin, Kurt Johansson, Patrick Ferrari, Jinho Baik, Kazumasa Takeuchi, Kostya Khanin Herbert Spohn and Craig Tracy for numerous illuminating discussions. An essential part of this work was done during the workshop "New approaches to non-equilibrium and random systems: KPZ integrability, universality, applications and experiments" (Jan 11 - Mar 11, 2016) at Kavli Institute of Theoretical Physics, University of California, Santa Barbara. This research was supported in part by the National Science Foundation under Grant No. NSF PHY11-25915.

Appendix

In the notations of the present paper the general result of the paper [33], (eqs(73), (76) and (79)-(82)) for the probability distribution function, eq.(14), is given by

$$W(f_1,f_2,\Delta) \;=\; \sum_{M_1=0}^{\infty}\frac{(-1)^{M_1}}{(M_1!)^2}\sum_{M_2=0}^{\infty}\frac{(-1)^{M_2}}{M_2!}\prod_{\alpha=1}^{M_1+M_2}\Big[\int_0^{\infty}du_\alpha\Big]\prod_{\beta=1}^{M_1}\Big[\int_0^{\infty}dv_\beta\Big]\times \tag{A.1}$$

$$\times\sum_{\mathcal{P}\in S_{M_1+M_2}}(-1)^{[\mathcal{P}]}\sum_{\tilde{\mathcal{P}}\in S_{M_1}}(-1)^{[\tilde{\mathcal{P}}]}\prod_{\beta=1}^{M_1}\Big[G\big(u_\beta,v_\beta;\;u_{\mathcal{P}_\beta},v_{\tilde{\mathcal{P}}_\beta}\big)\Big]\times$$

$$\times\prod_{\gamma=1}^{M_2}\Big[2^{1/3}K\big[2^{1/3}u_{M_1+\gamma}-\tilde{f}_2;\;2^{1/3}u_{\mathcal{P}_{M_1+\gamma}}-\tilde{f}_2\big]\Big]$$

where $\mathcal{P}$ and $\tilde{\mathcal{P}}$ denote the permutations of (M_1+M_2) variables $\{u_\alpha\}$ and M_1 variables $\{v_\beta\}$ correspondingly, $K(u;u')=\int_0^{+\infty}dy\;\mathrm{Ai}(y+u)\;\mathrm{Ai}(y+u')$ is the Airy kernel and

$$G\big(u,u';\;v,v'\big) \;=\; \sum_{i=1}^{4}G_i\big(u,u';\;v,v'\big) \tag{A.2}$$

with

$$G_1\big(u,u';\;v,v'\big) \;=\; 2^{2/3}\big(1-\Delta\big)^{2/3}\,\mathrm{Ai}\Big[2^{1/3}\big(1-\Delta\big)^{1/3}(u+v')-\tilde{f}_1\Big]\;\mathrm{Ai}\Big[2^{1/3}\big(1-\Delta\big)^{1/3}(u'+v)-\tilde{f}_1\Big] \tag{A.3}$$

$$G_2\big(u,u';\;v,v'\big) \;=\; -2^{2/3}\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}K\Big[2^{1/3}u-\tilde{f}_2;\;2^{1/3}u'-\tilde{f}_2\Big]\times$$

$$\times\;K\Bigg[2^{1/3}\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}v-\frac{\tilde{f}_1-(1-\Delta)^{1/3}\tilde{f}_2}{\Delta^{1/3}}\,;\;2^{1/3}\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}v'-\frac{\tilde{f}_1-(1-\Delta)^{1/3}\tilde{f}_2}{\Delta^{1/3}}\Bigg] \tag{A.4}$$

$$G_3\big(u,u';\;v,v'\big) \;=\; -2\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}\int_{-\infty}^{+\infty}\frac{dp}{2\pi}\int_0^{+\infty}dy\;\times$$

$$\times\;\mathrm{Ai}\Bigg[-\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}y+\Big(\frac{1-\Delta}{\Delta}\Big)^{-2/3}p^2-\frac{f_1-(1-\Delta)^{1/3}f_2}{\Delta^{1/3}}+\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}(v+v')\Bigg]\times$$

$$\times\;\mathrm{Ai}\big(y+p^2-f_2+u+u'\big)\exp\{ip\,(u-u'-v+v')\} \tag{A.5}$$

$$G_4\big(u,u';\,v,v'\big) \;=\; -4\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}\int\int_{-\infty}^{+\infty}\frac{dp\,dq}{(2\pi)^2}\int_0^{+\infty}dy\;\times$$
$$\times\;\mathrm{Ai}\bigg[-\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}y+\Big(\frac{1-\Delta}{\Delta}\Big)^{-2/3}p^2-\frac{f_1-(1-\Delta)^{1/3}f_2}{\Delta^{1/3}}+\Big(\frac{1-\Delta}{\Delta}\Big)^{1/3}(v+v')\bigg]\times$$
$$\times\;\mathrm{Ai}\big(y+q^2-f_2+u+u'\big)\frac{\sin\big[(q-p)\,y\big]}{(q-p)}\exp\{iq(u-u')-ip(v-v')\} \tag{A.6}$$

where $\tilde{f}_{1,2}\equiv 2^{-2/3}\,f_{1,2}$

Introducing the new scaling parameter ξ, eq.(17), and taking the limit $\Delta\to 0$ (such that the parameter ξ remains finite) we get:

$$G_1\big(u,u';\,v,v'\big)\Big|_{\Delta\to 0}\;\simeq\;2^{2/3}\,\mathrm{Ai}\Big[2^{1/3}(u+v')-\tilde{f}_1\Big]\;\mathrm{Ai}\Big[2^{1/3}(u'+v)-\tilde{f}_1\Big] \tag{A.7}$$

$$G_2\big(u,u';\,v,v'\big)\Big|_{\Delta\to 0}\;\simeq\;-\frac{2^{2/3}}{\Delta^{1/3}}K\Big[2^{1/3}u-\tilde{f}_2;\;2^{1/3}u'-\tilde{f}_2\Big]\;K\bigg[\frac{2^{1/3}v}{\Delta^{1/3}}-\xi\,;\;\frac{2^{1/3}v'}{\Delta^{1/3}}-\xi\bigg] \tag{A.8}$$

Redefining $y\to\Delta^{1/3}y$ in eq.(A.5) and taking the limit $\Delta\to 0$ we obtain

$$G_3\big(u,u';\,v,v'\big)\Big|_{\Delta\to 0}\;\simeq\;-2\int_{-\infty}^{+\infty}\frac{dp}{2\pi}\int_0^{+\infty}dy\;\times$$
$$\times\;\mathrm{Ai}\bigg(-y-\tilde{\xi}+\frac{v+v'}{\Delta^{1/3}}\bigg)\;\mathrm{Ai}\big(p^2-f_2+u+u'\big)\exp\{ip\,(u-u'-v+v')\} \tag{A.9}$$

Finally, redefining $y\to\Delta^{1/3}y$, $p\to\Delta^{-1/3}p$ in eq.(A.6) and taking the limit $\Delta\to 0$ we find

$$G_4\big(u,u';\,v,v'\big)\Big|_{\Delta\to 0}\;\simeq\;-4\int\int_{-\infty}^{+\infty}\frac{dp\,dq}{(2\pi)^2}\int_0^{+\infty}dy\;\frac{\sin\big(p\,y\big)}{p}\;\times$$
$$\times\;\mathrm{Ai}\bigg(-y+p^2-\tilde{\xi}+\frac{v+v'}{\Delta^{1/3}}\bigg)\;\mathrm{Ai}\big(q^2-f_2+u+u'\big)\exp\{iq(u-u')-ip(v-v')\} \tag{A.10}$$

Redefining $v_\beta\;\to\;\Delta^{1/3}\,2^{-1/3}\,v_\beta$ and $u_\alpha\;\to\;2^{-1/3}\,u_\alpha$ in eq.(A.1) we find that the kernel $G\big(u,u';\;\;v,v'\big)\;=\;\sum_{i=1}^{4}G_i\big(u,u';\,v,v'\big)$ gets the multiplicative factor $\Delta^{1/3}$. According to eqs.(A.7), (A.9) and (A.10), we find that

$$\Delta^{1/3}\,G_{1,3,4}\big(u,u';\,v,v'\big)\big|_{\Delta\to 0}\;\to\;0 \tag{A.11}$$

According to eq.(A.8), the only contribution which remains finite in the limit $\Delta\;\to\;0$ is:

$$\Delta^{1/3}\,G_2\big(u,u';\,v,v'\big)\;\to\;-K\big(u-\tilde{f}_2;\;u'-\tilde{f}_2\big)\;K\big(v-\xi\,;\;v'-\xi\,\big) \tag{A.12}$$

Thus, substituting $\tilde{f}_2=f,\;\;\tilde{f}_1\;=\;f\;+\;\Delta^{1/3}\xi,\;\;v_\beta\;\to\Delta^{1/3}\,2^{-1/3}\,v_\beta$ and $u_\alpha\to 2^{-1/3}\,u_\alpha$ in eq.(A.1), in the limit $\Delta\;\to\;0$ we get

$$
\begin{aligned}
W(f,\xi) \;&\equiv\; \lim_{\Delta\to 0} W\Big(2^{2/3}f \,+\, 2^{2/3}\Delta^{1/3}\,\xi,\;\; 2^{2/3}f;\;\Delta\Big) \\
&=\; \sum_{M_1=0}^{\infty}\frac{(-1)^{M_1}}{(M_1!)^2}\sum_{M_2=0}^{\infty}\frac{(-1)^{M_2}}{M_2!}\prod_{\alpha=1}^{M_1+M_2}\left[\int_0^{\infty}du_\alpha\right]\prod_{\beta=1}^{M_1}\left[\int_0^{\infty}dv_\beta\right]\sum_{\mathcal{P}\in S_{M_1+M_2}}(-1)^{[\mathcal{P}]}\sum_{\tilde{\mathcal{P}}\in S_{M_1}}(-1)^{[\tilde{\mathcal{P}}]}\;\times \\
&\times\;\prod_{\beta=1}^{M_1}\Big[-K\big(u_\beta-f;\;u_{\mathcal{P}_\beta}-f\big)\;K\big(v_\beta-\xi\,;\;v_{\tilde{\mathcal{P}}_\beta}-\xi\,\big)\Big]\;\prod_{\gamma=1}^{M_2}\Big[K\big(u_{M_1+\gamma}-f;\;u_{\mathcal{P}_{M_1+\gamma}}-f\big)\Big] \\
&=\; \sum_{M_1=0}^{\infty}\frac{1}{(M_1!)^2}\sum_{M_2=0}^{\infty}\frac{(-1)^{M_2}}{M_2!}\prod_{\alpha=1}^{M_1+M_2}\left[\int_0^{\infty}du_\alpha\right]\prod_{\beta=1}^{M_1}\left[\int_0^{\infty}dv_\beta\right]\times \\
&\times\;\sum_{\mathcal{P}\in S_{M_1+M_2}}(-1)^{[\mathcal{P}]}\prod_{\alpha=1}^{M_1+M_2}\Big[K\big(u_\alpha-f;\;u_{\mathcal{P}_\alpha}-f\big)\Big]\sum_{\tilde{\mathcal{P}}\in S_{M_1}}(-1)^{[\tilde{\mathcal{P}}]}\prod_{\beta=1}^{M_1}\Big[K\big(v_\beta-\xi\,;\;v_{\tilde{\mathcal{P}}_\beta}-\xi\,\big)\Big] \qquad \text{(A.13)}
\end{aligned}
$$

or

$$
\begin{aligned}
W(f,\xi) \;&=\; \sum_{M=0}^{\infty}\frac{(-1)^{M}}{M!}\prod_{\alpha=1}^{M}\left[\int_0^{\infty}du_\alpha\right]\det\Big(K\big[u_\alpha-f;\;u_{\alpha'}-f\big]\Big)\Big|_{(\alpha,\alpha')=1,\ldots,M}\;\times \\
&\times\;\sum_{M_1=0}^{M}\frac{(-1)^{M_1}M!}{(M_1!)^2(M-M_1)!}\prod_{\beta=1}^{M_1}\left[\int_0^{\infty}dv_\beta\right]\det\Big(K\big[v_\beta-\xi;\;v_{\beta'}-\xi\big]\Big)\Big|_{(\beta,\beta')=1,\ldots,M_1} \qquad \text{(A.14)}
\end{aligned}
$$

It can be easily shown (see below) that the above formula can be represented in the compact form in terms of the product of two Fredholm determinants [45]

$$
W(f,\xi) \;=\; \frac{1}{2\pi i}\int_{|z|=1}\frac{dz}{z}\det\Big[\hat{1}\;-\;(1-z)\hat{K}_{-f}\Big]\;\det\Big[\hat{1}\;+\;z^{-1}\hat{K}_{-\xi}\Big] \qquad \text{(A.15)}
$$

where the integration over z goes in the complex plane around zero and $\hat{K}_{-f}$ is the integral operator with the usual Airy kernel $K(u-f;u'-f)$ (with $(u,\,u')\;\geq 0$). Indeed, expanding the Fredholm determinants in eq.(A.15), one gets

$$
W(f,\xi) \;=\; \frac{1}{2\pi i}\int_{|z|=1}\frac{dz}{z}\sum_{M=0}^{\infty}\frac{(-1)^{M}}{M!}\,(1-z)^{M}D_M(-f)\;\sum_{M_1=0}^{\infty}\frac{1}{M_1!}\,z^{-M_1}D_{M_1}(-\xi) \qquad \text{(A.16)}
$$

where

$$
D_M(s) \;\equiv\; \prod_{\alpha=1}^{M}\left[\int_0^{\infty}du_\alpha\right]\det\Big(K\big[u_\alpha+s;\;u_{\alpha'}+s\big]\Big)\Big|_{(\alpha,\alpha')=1,\ldots,M} \qquad \text{(A.17)}
$$

Simple algebra yields:

$$
\begin{aligned}
W(f,\xi) \;&=\; \sum_{M=0}^{\infty}\frac{(-1)^{M}}{M!}D_M(-f)\sum_{M_1=0}^{\infty}\frac{1}{M_1!}D_{M_1}(-\xi)\sum_{L=0}^{M}\frac{(-1)^{L}M!}{L!(M-L)!}\;\frac{1}{2\pi i}\int_{|z|=1}\frac{dz}{z}\;z^{L-M_1} \\
&=\; \sum_{M=0}^{\infty}\frac{(-1)^{M}}{M!}D_M(-f)\sum_{M_1=0}^{\infty}\frac{1}{M_1!}D_{M_1}(-\xi)\sum_{L=0}^{M}\frac{(-1)^{L}M!}{L!(M-L)!}\;\delta_{L,M_1} \qquad \text{(A.18)}
\end{aligned}
$$

which coinsides with eq.(A.14).

[1] Kardar M, Parisi G, Zhang Y-C (1986) Phys. Rev. Lett. **56**, 889
[2] Halpin-Healy T and Zhang Y-C (1995) Phys. Rep. **254**, 215
[3] Burgers J M (1974) *The Nonlinear Diffusion Equation* (Reidel, Dordrecht)
[4] Kardar M (2007), *Statistical physics of fields* (Cambridge: Cambridge University Press)
[5] Huse D A , Henley C L and Fisher D S (1985) Phys. Rev. Lett. **55**, 2924
[6] Huse D A and Henley C L (1985) Phys. Rev. Lett. **54**, 2708
[7] Kardar M and Zhang Y-C (1987) Phys. Rev. Lett. **58**, 2087
[8] Kardar M (1987) Nucl. Phys. **B 290**, 582
[9] Bouchaud J P and Orland H (1990) J. Stat. Phys. **61**, 877
[10] Brunet E and Derrida B (2000) Phys. Rev. E **61**, 6789
[11] Johansson K (2000) Comm. Math. Phys. **209**, 437
[12] Prahofer M and Spohn H (2002) J. Stat. Phys. **108**, 1071
[13] Ferrari P L and Spohn H (2006) Comm. Math. Phys. **265**, 1
[14] Sasamoto T and Spohn H (2010) Phys. Rev. Lett. **104**, 230602
[15] Sasamoto T and Spohn H (2010) Nucl. Phys. **B834**, 523
[16] Sasamoto T and Spohn H (2010) J. Stat. Phys. **140**, 209
[17] Amir G , Corwin I and Quastel J (2011) Comm. Pure Appl. Math. **64**, 466
[18] Dotsenko V and Klumov B (2010) J.Stat.Mech. P03022
[19] Dotsenko V (2010) EPL, **90**, 20003
[20] Dotsenko V (2013) J.Stat.Mech. P07010
[21] Calabrese P, Le Doussal P and Rosso A (2010) EPL, **90**, 20002
[22] Calabrese P and Le Doussal P (2011) Phys. Rev. Lett. **106**, 250603
[23] Dotsenko V (2012) J. Stat. Mech. P11014
[24] Gueudré T and Le Doussal P (2012) EPL, **100**, 26006
[25] Corwin I (2012) Random Matrices: Theory Appl. **1**, 1130001
[26] Borodin A, Corwin I and Ferrari P (2012) arXiv:1204.1024
[27] Tracy C A and Widom H (1994) Commun.Math Phys. **159**, 151
[28] Prolhac S and Spohn H (2011) J.Stat.Mech. P01031
[29] Dotsenko V (2013) J.Phys. A: Math. Theor. bf 46, 355001
[30] Imamura T, Sasamoto T and Spohn H (2013) J.Phys. A: Math. Theor. bf 46, 355002
[31] Prolhac S and Spohn H (2011) J.Stat.Mech. P03020
[32] Dotsenko V (2014), Cond.Mat.Phys. **17**, 33003
[33] Dotsenko V (2013) J.Stat.Mech. P06017
[34] Dotsenko V (2015) J.Phys. A: Math. Theor. **48** 495001
[35] Johansson K (2015) arXiv:1502.00941
[36] Ferrari P L and Spohn H, *On time correlations for KPZ growth in one dimension*
[37] Takeuchi K A and Sano M (2012) J. Stat. Phys. **147**, 853-890
[38] Lieb E H and Liniger W (1963) Phys. Rev. **130**, 1605
[39] McGuire J B (1964) J. Math. Phys. **5**, 622
[40] Yang C N (1968) Phys. Rev. **168**, 1920
[41] Calabrese P and Caux J S (2007) Phys. Rev. Lett. **98**, 150403
[42] Dotsenko V (2011) *Universal Randomness*, Physics-Uspekhi, **54**(3), 259
[43] Dotsenko V (2012) J. Stat. Mech. P02012
[44] Gaudin M (1983) *La fonction d'onde de Bethe*, (Paris, Masson)
[45] I am grateful to Alexei Borodin for his indication that eq.(A.14) can be represented in the form of eq.(A.15)